

Extreme Energy ν_τ Propagation Through the Earth

F. Becattini, S. Bottai ¹

*University of Florence and INFN Sezione di Firenze
Largo E. Fermi 2, I-50125
Firenze (Italy)*

Abstract

The propagation of extremely energetic ν_τ 's through the Earth is studied by means of a detailed Monte Carlo simulation. All major mechanisms of ν_τ interactions and τ energy loss as well as all its relevant decay modes are properly taken into account. The detection possibility of ν_τ 's emerging from Earth in forthcoming neutrino telescopes is addressed.

To be published in Astroparticle Physics

¹e-mail: becattini@fi.infn.it, bottai@fi.infn.it

1 Introduction

Cosmic neutrinos with energy $> 10^{12}$ eV are a new and exciting observational window for cosmic ray physics and astrophysics. Indeed, many interesting models predict the existence of cosmic neutrinos with an energy up to $10^{18} - 10^{20}$ eV (*extreme energy*), where charged particles are expected to be cut-off by the so called GZK mechanism [1]. The aforementioned threshold of 10^{12} eV is necessary for a detection of diffuse flux, because of the large atmospheric neutrino background.

The detection of such extreme energy neutrinos is challenging as it demands a very large sensitive area. The experiments of the upcoming generation plan to use underwater-underice Cerenkov detectors [2] and large field-of-view atmospheric fluorescence detectors [3]. One of the best evidence of cosmic neutrinos would be the detection of upstream showers/particles emerging from the Earth. For this kind of events, the atmospheric muon and primary charged cosmic ray background would be completely suppressed. On the other hand, this signature can hardly be observed at extreme energy because the rise of weak cross sections entails the opacity of the Earth with respect to neutrino propagation [4, 5].

The problem of neutrino propagation through the Earth can be understood on the basis of neutrino interactions with matter. For neutrino energy well above few GeV, interactions with conventional matter are described by neutrino-nucleon deep inelastic scattering and neutrino-electron scattering [5]. The former can be classified in two main categories: charged current (CC) $\nu_l N \rightarrow l + \text{hadrons}$, where the neutrino disappears and a charged lepton l is created, and neutral current (NC) $\nu_l N \rightarrow \nu_l + \text{hadrons}$, where the neutrino survives the interaction. The neutrino-electron scattering cross section is several order of magnitude less than neutrino-nucleon cross section and does not affect significantly neutrino propagation in matter. The only exception is the W^- resonant scattering for the reaction $\bar{\nu}_e e \rightarrow W^- + \text{anything}$, which, however, hardly affects neutrino attenuation length in matter for $\bar{\nu}_e$ energy in the range $2 \cdot 10^{15} \text{ eV} \leq E_\nu \leq 2 \cdot 10^{16} \text{ eV}$ [5].

The neutrino-nucleon total cross section is approximately $\sigma_{tot} \approx 10^{-37} \text{ cm}^2$ at $E_\nu = 10 \text{ GeV}$ and rises roughly linearly with energy. As a consequence, while in the GeV energy range the Earth is completely transparent to neutrinos, it is expected to become opaque for sufficiently high energy. As to muon neutrinos, the CC interaction length inside Earth equals Earth's diameter for energy around 40 TeV [5]. In order to accurately describe the change in energy spectrum of a muon neutrino flux traversing the Earth, both the effect of neutrino-nucleon CC and NC must be properly taken into account. The problem of neutrino propagation in the Earth can be solved accurately either by Monte-Carlo simulation or by approximate iterative solution of the relevant transport equation [4].

It has been recently pointed out [6, 7, 8, 9] that the behaviour of τ -neutrinos, whose existence should be guaranteed in a neutrino-oscillation scenario, should be significantly different from ν_μ and ν_e . Whilst muon and electron neutrinos are practically absorbed after one CC interaction, the τ lepton created by the ν_τ CC scattering may decay in flight before losing too much energy, thereby generating a new ν_τ with comparable energy. Hence, ultra high energy τ -neutrinos should emerge from the Earth instead of being absorbed. For a correct evaluation of energy spectrum of ν_τ 's emerging from the Earth, one has to properly take into account ν_τ interactions as well as τ energy loss and decay.

An analytical approach similar to that used in [4] for ν_μ has been proposed by S. Iyer et al. [9] neglecting τ energy loss. This approach holds as long as energy does not exceed 10^{16} eV because above this energy τ interaction length becomes comparable with τ decay length (see next section).

In the present work, a detailed Monte Carlo calculation of $\nu_\tau - \tau$ system propagation through the Earth has been performed for energy up to 10^{20} eV including the τ energy loss contribution, with special emphasis on the initial ν_τ spectrum deformation as a function of the zenith angle of the emerging particle.

2 Neutrino cross sections and charged lepton interactions

The Montecarlo simulation has been performed following neutrinos and charged leptons along their path inside the Earth. The Earth model considered is the preliminary Earth model of ref. [10]. During particle propagation, the occurrence of interactions relevant for energy loss, decay or leptons production has been simulated by a unidimensional approach. The only secondary particles which have been followed are the ν_τ 's produced by τ decays and τ 's arising from ν_τ charged current interactions.

Deep inelastic neutrino-nucleon scattering is the dominant interaction of energetic neutrinos into conventional matter. Charged and neutral current differential cross sections used in this work have been calculated in the framework of QCD improved parton model. Charged current cross section for a neutrino-isoscalar nucleon interaction reads:

$$\frac{d^2\sigma}{dx dy} = \frac{2G_F^2 M E_\nu}{\pi} \left(\frac{M_W^2}{Q^2 + M_W^2} \right)^2 \left[xq(x, Q^2) + x\bar{q}(x, Q^2)(1 - y)^2 \right], \quad (1)$$

where $-Q^2$ is the invariant momentum transfer between the incident neutrino and outgoing muon, $\nu = E_\nu - E_l$ is the energy loss in the lab (target) frame, $x = Q^2/2M\nu$ and $y = \nu/E_\nu$, M and M_W are the nucleon and intermediate-boson masses, G_F is the Fermi constant and the quark distribution functions are:

$$\begin{aligned} q(x, Q^2) &= \frac{u_v(x, Q^2) + d_v(x, Q^2)}{2} + \frac{u_s(x, Q^2) + d_s(x, Q^2)}{2} + s_s(x, Q^2) + b_s(x, Q^2) \\ \bar{q}(x, Q^2) &= \frac{u_s(x, Q^2) + d_s(x, Q^2)}{2} + c_s(x, Q^2) + t_s(x, Q^2), \end{aligned} \quad (2)$$

where the subscripts v and s label valence and sea contributions, and u, d, c, s, t, b denote the distributions for various quark flavors.

Since a large contribution to cross sections comes from the very low x region laying beyond present accelerator domain, an extrapolation of parton distribution at very low x ($x \geq 10^{-8}$) is necessary. We have used the parton distribution set CTEQ3-DIS [11], available in the program library PDFLIB [12], with NLO-DGLAP formalism used for Q^2 evolution and assuming for very low x the same functional form measured at $x = \mathcal{O}(10^{-5})$. Even if more sophisticated approaches for low x extrapolation have been developed using

dynamical QCD [13] the results for cross section calculations (see Fig. 4) do not differ more than 10% from the approach taken here with CTEQ3-DIS plus "brute force" extrapolation [5, 13].

The electromagnetic interactions of muons with matter, at the actually considered energies, are dominated by radiative processes rather than ionisation [14]. The cross sections for electromagnetic radiative processes of τ are lower than muon's, still radiative interactions remain the dominant process for τ energy loss. The cross sections used for radiative electromagnetic interactions of τ leptons are based on QED calculation for Bremsstrahlung [15], for direct pair production [16] and for photonuclear interactions [17] by replacing muon or electron mass with τ mass (only formulae with explicit lepton mass dependence in the original paper have been considered). For all above processes we have implemented stochastic interactions for $\nu = (E_f - E_i)/E_i \geq 10^{-3}$ and a continuous energy loss for $\nu = (E_f - E_i)/E_i \leq 10^{-3}$, where E_i and E_f are the τ energies before and after the interaction respectively.

It is worth mentioning that Bremsstrahlung cross section scales as the inverse square of lepton mass whereas the direct pair production approximately scales according to m_e/m_l [18]. As a consequence, the dominant process of τ lepton energy loss is direct pair production rather than photon radiation. Along with electromagnetic interactions, we also considered weak interactions of τ with nucleons although they are almost negligible even at $E_\tau \sim 10^{20}$ eV. The τ decay has been simulated by using the TAUOLA package [19].

3 Simulation results

In Figs. 2, 3 scatter plots of initial versus final energy are shown. The initial energy is the energy of the neutrino before crossing the Earth while the final energy is the energy of neutrino emerging from the Earth. The initial energy distribution has been chosen to be proportional to E^{-1} so to have a uniform distribution in $\log E$. Only muon neutrinos either interacting with neutral current or not interacting at all are able to cross the Earth whereas the rest is absorbed in that the muon arising from a charged current interaction has a radiation length much lower than its decay length at any considered energy.

On the other hand, the τ generated from a ν_τ charged current interaction loses energy until its decay length becomes comparable to its radiation length, thereafter it decays into a new ν_τ . Unlike for muons, this occurs around $10^{16} - 10^{17}$ eV and, as a consequence, CC interacting ν_τ 's with $E_\tau \geq 10^{17}$ eV, instead of being absorbed, populate the $E_\tau \leq 10^{17}$ eV energy region. The result of our Monte Carlo calculation shown in Fig. 3 is qualitatively in a fairly good agreement with a similar plot in ref. [7], though a more quantitative comparison could be desirable. The ν_τ propagation develops in several $\nu_\tau - \tau$ transformation steps. The accumulation of ν_τ energies below 10^{17} eV is modulated by the zenith angle of the emerging particle because particle traverse different amounts of matter. In Fig. 4 a scatter plot of final energy versus θ_{zenith} is shown. The input energy spectrum has been chosen once again $\propto E^{-1}$ while the zenith angle distribution is uniform in order to better display of the final energy angular dependence. The final energies cluster around $0.5 \cdot 10^{17}$ eV for nearly horizontal particles and go down to 10^{14} eV for vertically incoming

particles.

We have studied the final energy spectrum of tau and muon neutrinos crossing the Earth at three different incident zenith angles for three different initial energy spectra: $\propto E^{-1}$ (see Fig. 5), $\propto E^{-2}$ (see Fig. 6) and a possible energy spectrum predicted for topological defect decays [20] (see Fig. 7). The main feature of ν_τ spectrum is indeed the presence of a characteristic bump before a steep drop, due to the ν_τ regeneration mechanism described above. This clearly shows up for hard initial spectra (see Fig. 5) whereas it is hardly recognizable for softer initial spectra (see Fig. 7) and becomes a smooth slope in case of steeply falling energy spectra (see Fig. 6). The bump position strongly depends on zenith angle because energy loss increases as a function of encountered matter thickness. The results obtained in this work are in good agreement with those by S. Iyer et al. [9], who took an analytical approach. The main difference between the two results is an excess of ν_τ 's around the characteristic bump in the $\propto E^{-1}$ energy spectrum in our work. This can be easily explained by taking into account the difference in the initial neutrino spectrum. We have used a pure $\propto E^{-1}$ spectrum up to 10^{20} eV while S. Iyer et al. used a modulated cutoff for energies greater than 10^{17} eV. Since in a $\propto E^{-1}$ spectrum the number of neutrinos which are shifted from high energy to the characteristic bump at lower energy receives approximately an equal contribution for each energy decade above the bump, the agreement is restored. Furthermore, in ref. [9], τ energy loss is not taken into account. However, the effect of such approximation is almost negligible because in the energy region considered therein, below 10^{15} eV, τ decay length is much less than its interaction length.

4 Conclusions

Whilst, as it is well known, the Earth start to be opaque for muon neutrino propagation at energies around 10 TeV, a regeneration mechanism relevant to ν_τ prevents them from being absorbed. This mechanism is ultimately owing to the shorter, with respect to muon, τ lepton decay length. Nevertheless, the unavoidable radiative τ energy loss sets an upper bound on the energy of emerging ν_τ 's to 10^{17} eV. Therefore, for underwater-underice neutrino telescopes, whose sensitivity will cover the whole neutrino spectrum, ν_τ 's could be the major source of neutrino events above the PeV region. In such detectors the characteristics of Fig. 5 could be also used as one of the signatures of tau neutrinos. For atmospheric fluorescence detectors the effective cut-off of 10^{17} eV could be below their energy threshold. If this is the case, those detectors should be able to detect only almost horizontal neutrinos.

Acknowledgments

We are very grateful to V. Naumov for bringing our attention to the problem of ν_τ propagation and for very fruitful discussions.

References

- [1] K. Greisen, Phys. Rev. Lett. 16 (1996) 748; G.T. Zatsepin and V.A. Kuzmin, JETP Lett. 4 (1996) 78
- [2] T.K. Gaisser, F. Halzen, T. Stanev, Phys. Rep. 258 (1995) 173 and references therein
- [3] J. Linsley, L. Scarsi, P. Spillantini, Y. Takahashi in Proc. of 25th ICRC vol. 5 (1997) 389
- [4] V. Naumov, L. Perrone, Astropart. Phys. 10 (1999) 239 and references therein
- [5] Gandhi et al., Astropart. Phys. 5 (1996) 81
- [6] D. Fargion, "The Role of Tau Neutrino Ultrahigh-energy astrophysics in km^3 detectors", astro-ph/9704205 (1997)
- [7] F. Halzen, D. Saltzberg, Phys. Rev. Lett. 81 (1998) 4308
- [8] F. Becattini, S. Bottai, 26th ICRC, vol. 2 (1999) 249
- [9] S. Iyer et al., Phys. Rev. D61 (2000) 053003
- [10] A.M. Dziewonski and D.L. Anderson, Physics Earth and Planetary Interiors, 25 (1981) 297
- [11] H. Lai et al., Phys. Rev. D51 (1995) 4763
- [12] H. Plothow-Besch, "PDFLIB, Version 7.09", CERN-PPE report 1997.07.02
- [13] M. Gluck, S.Kretzer and E. Reya, Astropart. Phys. 11 (1999) 327
- [14] P. Lipari, T. Stanev, Phys. Rev. D44 (1991) 3543
- [15] A.A. Petrukhin, V.V. Shestakov, Can. J. Nucl. Phys, 46 (1968) S377.
- [16] R.P. Kokoulin and A.A. Petrukhin, Acta Phys. Hung. 29 Suppl. 4 (1970) 277
- [17] L.B. Bezrukov and E.V. Bugaev Sov. J.Nucl. Phys. 33 (1981) 635
- [18] M.J. Tannenbaum, Nucl. Inst. Meth. A300 (1991) 595
- [19] S. Jadach, Z. Was, R. Decker, J. H. Kuhn, CERN-TH 6793/93 (1993)
- [20] G. Sigl, S. Lee, D. N. Schramm and P. Coppi, Phys. Lett. B392 (1997) 129




Figure 1: Total ν_τ CC and NC cross sections on isoscalar nucleon target as a function of energy.




Figure 2: Scatter plot of initial versus final energy of ν_μ propagating through the Earth with an incident zenith angle uniformly distributed in the range $95^\circ \leq \theta_{\text{zenith}} \leq 180^\circ$.




Figure 3: Scatter plot of initial versus final energy of ν_τ propagating through the Earth with an incident zenith angle uniformly distributed in the range $95^\circ \leq \theta_{zenith} \leq 180^\circ$.




Figure 4: Scatter plot of final energy as a function of zenith angle for an initial energy spectrum $\propto E^{-1}$.




Figure 5: Final energy spectra of ν_τ 's and ν_μ 's crossing the Earth at zenith angles of 180° , 140° and 100° for an initial energy spectrum $\propto E^{-1}$.




Figure 6: Final energy spectra of ν_τ 's and ν_μ 's crossing the Earth at zeinth angles of 180° , 140° and 100° for an initial energy spectrum $\propto E^{-2}$.




Figure 7: Final energy spectra of ν_{τ} 's and ν_{μ} 's crossing the Earth at zenith angles of 180° , 140° and 100° for an initial energy spectrum calculated for topological defect decays in ref. [20].